

CORRECTION

In the article entitled “**Normative values of the balloon expulsion test in healthy adults**” by Dedeli et al (**Turk J Gastroenterol 2007; 18: 177-181**) the following corrections should be noted.

Data under Table 2 should be listed under Table 3 and data under Table 3 should be listed under Table 2 (page 179). We regret the error and provide you below the original and revised versions of tables 2 and 3. The tables have been also corrected on the Journal's Web site at www.turkgastro.org.

Original Version

Table 2. Balloon expulsion times according to age subgroups in men

Age group	Mean (sec)	N	Std. Deviation	Minimum	Maximum	Mean Rank
20-39	58.6	22	25.9	20	90	32.863
40-59	54.7	23	20.1	20	90	29.369
60+	55.9	16	16.8	17	90	30.781
Total	56.4	61	21.3	17	90	

sec: Second, std: Standard

Table 3. Balloon expulsion times according to age subgroups in women

Age group	Mean (sec)	n	Std. Deviation	Minimum	Maximum	Mean Rank
20-39	28.6	20	11.7	15	60	14.625
40-59	49.8	20	19.5	20	85	32.975
60+	57.1	15	11.2	35	83	39.2
Total	44.1	55	19.0	15	85	

sec: Second, std: Standard

Revised Version

Table 2. Balloon expulsion times according to age subgroups in men

Age group	Mean (sec)	n	Std. Deviation	Minimum	Maximum	Mean Rank
20-39	28.6	20	11.7	15	60	14.625
40-59	49.8	20	19.5	20	85	32.975
60+	57.1	15	11.2	35	83	39.2
Total	44.1	55	19.0	15	85	

sec: Second, std: Standard

Table 3. Balloon expulsion times according to age subgroups in women

Age group	Mean (sec)	N	Std. Deviation	Minimum	Maximum	Mean Rank
20-39	58.6	22	25.9	20	90	32.863
40-59	54.7	23	20.1	20	90	29.369
60+	55.9	16	16.8	17	90	30.781
Total	56.4	61	21.3	17	90	

sec: Second, std: Standard